

**INTERNATIONAL ECOLOGICAL
CLASSIFICATION STANDARD:**

TERRESTRIAL ECOLOGICAL CLASSIFICATIONS

Ecological Systems of Texas' Cross Timbers

08 October 2009

by

NatureServe

1101 Wilson Blvd., 15th floor
Arlington, VA 22209

This subset of the International Ecological Classification Standard covers terrestrial ecological systems attributed to the Texas. This classification has been developed in consultation with many individuals and agencies and incorporates information from a variety of publications and other classifications. Comments and suggestions regarding the contents of this subset should be directed to [Judy Teague <judy_teague@natureserve.org>].



Copyright © 2009 NatureServe, 1101 Wilson Blvd, 15th floor
Arlington, VA 22209, U.S.A. All Rights Reserved.

Citations:

The following citation should be used in any published materials which reference ecological system and/or International Vegetation Classification (IVC hierarchy) and association data:

NatureServe. 2009. International Ecological Classification Standard: Terrestrial Ecological Classifications. NatureServe Central Databases. Arlington, VA. U.S.A. Data current as of 08 October 2009.

TABLE OF CONTENTS

FOREST AND WOODLAND..... 3
 CES205.682 Crosstimbers Oak Forest and Woodland 3
 CES303.656 Edwards Plateau Dry-Mesic Slope Forest and Woodland 3
 CES303.660 Edwards Plateau Limestone Savanna and Woodland 3
 CES303.038 Edwards Plateau Mesic Canyon 3

SHRUBLAND 5
 CES303.041 Edwards Plateau Limestone Shrubland..... 5
 CES303.671 Western Great Plains Sandhill Steppe 5

HERBACEOUS 6
 CES303.659 Central Mixedgrass Prairie..... 6
 CES205.685 Southeastern Great Plains Tallgrass Prairie 6
 CES205.684 Texas Blackland Tallgrass Prairie..... 7
 CES303.670 Western Great Plains Sand Prairie 7

WOODY WETLAND 9
 CES303.651 Edwards Plateau Floodplain 9
 CES303.652 Edwards Plateau Riparian 9
 CES205.710 Southeastern Great Plains Floodplain Forest..... 9
 CES205.709 Southeastern Great Plains Riparian Forest..... 9

BARREN 11
 CES203.398 Southeastern Coastal Plain Cliff..... 11

FOREST AND WOODLAND

CES205.682 CROSTIMBERS OAK FOREST AND WOODLAND

This system is primarily found within central Texas and Oklahoma, ranging north to southeastern Kansas, and east into eastern Oklahoma. It is distinct from the surrounding prairie by the higher density of tree species. The area consists of irregular plains with primarily sandy to loamy Ustalf soils that range from shallow to moderately deep. Rainfall can be moderate, but somewhat erratic, therefore moisture is often limiting during part of the growing season. Short, stunted *Quercus stellata* and *Quercus marilandica* characterize and dominate this system. Other species, such as *Carya texana*, *Carya cordiformis*, *Quercus prinoides*, *Ulmus crassifolia*, and *Quercus* spp., can also be present within their respective ranges. The understory often contains species typical of the surrounding prairies, in particular *Schizachyrium scoparium*. Shrubs such as *Rhus* spp. may also be present. Drought, grazing, and fire are the primary natural processes that affect this system. Overgrazing and conversion to agriculture, along with fire suppression, have led to the invasion of some areas by problematic brush species such as *Juniperus virginiana* and *Juniperus ashei* and *Prosopis glandulosa* farther south in Texas and Oklahoma. It has also led to decreases in native grass cover allowing for annual grasses and forbs to invade.

This system is located on irregular plains comprised of sandy to loamy Ustalf soils. These soils range from shallow to moderately deep. Rainfall can be moderate, but sporadic, leading to periods of limiting moisture.

CES303.656 EDWARDS PLATEAU DRY-MESIC SLOPE FOREST AND WOODLAND

This system occurs on dry-mesic, middle slopes of the rolling uplands of the Edwards Plateau of Texas. The canopy is typically dominated by deciduous trees, including *Quercus buckleyi*, *Fraxinus texensis*, or *Ulmus crassifolia*. *Quercus fusiformis* and *Juniperus ashei* are often present but not dominant in this system. Canopy closure is variable, and this system can be expressed as forests and woodlands. This system occurs on dry-mesic, primarily north- and east-facing limestone slopes in the Edwards Plateau of Texas.

CES303.660 EDWARDS PLATEAU LIMESTONE SAVANNA AND WOODLAND

This upland system occurs primarily on limestone soils in the Edwards Plateau and forms the matrix within this ecoregion. It can also occur on limestone in the shortgrass regions of Texas and north into Oklahoma in areas such as the Arbuckle Mountains. This system is typified by a mosaic of evergreen oak forests, woodlands and savannas over shallow soils of rolling uplands and upper slopes within the Edwards Plateau and Lampasas Cutplain. *Quercus fusiformis* or *Juniperus ashei* typically dominate the canopy of this system. Other species may include *Quercus buckleyi*, *Quercus laceyi*, *Quercus stellata*, *Ulmus crassifolia*, *Fraxinus texensis*, *Quercus sinuata*, *Quercus vaseyana*, and *Diospyros texana*. Physiographic expression of this system varies from dense mottes (patches of forest where canopy cover approaches 100%) interspersed with grasslands to open savannalike woodlands with scattered individual or small groups of trees. Understories can contain various shrubs and graminoids, including *Cercis canadensis* var. *texensis*, *Forestiera pubescens*, *Sideroxylon lanuginosum*, *Diospyros texana*, *Rhus trilobata*, *Bouteloua* spp., *Schizachyrium scoparium*, *Nassella leucotricha*, *Carex planostachys*, *Aristida purpurea*, *Aristida oligantha*, *Liatris mucronata*, *Stillingia texana*, *Symphotrichum ericoides*, *Hedyotis nigricans*, *Monarda citriodora*, and *Salvia texana*. Grasslands dominated by *Schizachyrium scoparium* occur in small patches within more closed woodlands and in larger patches between mottes or in open savannalike woodlands with scattered trees. Grasslands in this system tend to grade from shortgrass communities in the west to mixedgrass communities to the east. Substrate (limestone) determines the range of this system within given examples. Some disturbed areas of the western plateau are now dominated by mesquite woodland. Natural mesquite woodlands are believed to have occurred on the deeper soils of adjacent riparian systems. This system is primarily restricted to limestone soils of rolling uplands within the Cretaceous limestone formations of the Edwards Plateau and dissected Pennsylvanian limestone formations within Texas and north into Oklahoma. Soil moisture and topography influence this system.

CES303.038 EDWARDS PLATEAU MESIC CANYON

This system is largely endemic to the Edwards Plateau ecoregion and occurs on canyon bottoms, mesic lower slopes and steep canyons, primarily in the Southern Balcones Escarpment, but also in the Eastern Balcones Escarpment. This system also includes cliff faces and lower slopes of boxed canyons occurring as narrow, sometimes long bands in areas often with seeps where moisture is consistently more available than on adjacent slopes. The tree canopy is generally closed. Common components include *Ulmus crassifolia*, *Juglans major*, *Quercus buckleyi*, *Quercus laceyi*, *Prunus serotina* var. *eximia* (becoming less common to the north), *Fraxinus texensis* (dominant in the northeastern plateau), *Quercus muehlenbergii*, and *Acer grandidentatum*. Canyon bottoms may have scattered *Quercus macrocarpa*. Substrate (limestone) and topographic position (north and east aspects and lower slopes) are the

dominant characteristics of this system. Small seepage areas are often dominated by *Adiantum capillus-veneris*, with *Thelypteris ovata* var. *lindheimeri* on nearby moist habitats. Other prominent species include *Buddleja racemosa*, *Ungnadia speciosa*, and *Toxicodendron radicans* ssp. *eximium*. Fire probably plays little role in the system, while grazing and browsing (by native as well as exotic ungulates) may play an important role in recruitment and understory composition. Adjacent, drier slopes are usually dominated by various *Quercus* species and *Juniperus ashei*. Largely endemic to the Edwards Plateau ecoregion and occurs on canyon bottoms, mesic lower slopes and steep canyons, primarily in the Southern Balcones Escarpment, but also in the Eastern Balcones Escarpment. This system occurs on mesic lower slopes, primarily with northern and eastern aspects and steep canyons over limestone in the Edwards Plateau region of Texas.

SHRUBLAND

CES303.041 EDWARDS PLATEAU LIMESTONE SHRUBLAND

This ecological system occurs as a matrix on relatively thin-soiled surfaces of plateaus of the massive limestones such as the Edwards limestone. These short to tall shrublands are variable in density depending on the relative amount of, and depth to, bedrock. *Quercus sinuata* var. *breviloba* is an important component of the system, with some areas dominated by *Quercus fusiformis*. *Juniperus ashei* is often an important component of this system. In the west, *Pinus remota* may also contribute to a scattered emergent overstory. Other shrub species may include *Rhus virens*, *Rhus lanceolata*, *Cercis canadensis* var. *texensis*, *Forestiera pubescens*, *Forestiera reticulata*, *Fraxinus texensis*, *Ungnadia speciosa*, *Sophora secundiflora*, *Diospyros texana*, *Salvia ballotiflora*, *Mimosa borealis*, *Condalia hookeri*, *Rhus trilobata*, *Opuntia engelmannii*, and *Mahonia trifoliolata*. This system also includes *Quercus mohriana*- or *Quercus vaseyana*-dominated shrublands that are more common to the west, often sharing dominance with *Juniperus pinchotii*. Herbaceous cover may be patchy and is generally graminoid with species including *Schizachyrium scoparium*, *Bouteloua curtipendula*, *Bouteloua rigidisetata*, *Bouteloua trifida*, *Hilaria belangeri*, *Bothriochloa laguroides* ssp. *torreyana*, *Nassella leucotricha*, *Erioneuron pilosum*, *Aristida* spp., and others. Disturbances such as fire may be important processes maintaining this system. However, it appears to persist on thin-soiled sites. In the western portions of the Edwards Plateau, more xeric conditions lead to the slow succession of sites to woodlands, resulting in long-persisting shrublands. This system occurs on thin soils over limestone in the Edwards Plateau of Texas.

This system occurs in a steady state on thin-soiled xeric sites. Shrub cover can be 100% in patches, but overall cover may be 40-50%. Patches of dense shrubs may be interspersed with bare rock and grasslands over shallow soil. Farther west this system grades into other shallow-soiled shrubland systems.

CES303.671 WESTERN GREAT PLAINS SANDHILL STEPPE

This system is found mostly in south-central areas of the Western Great Plains Division ranging from southwestern Wyoming and southwestern Nebraska up into the Nebraska Sandhill region, south through eastern Colorado, and New Mexico to central Texas, although some examples may reach as far north as the Badlands of South Dakota. The climate is semi-arid to arid for much of the region in which this system occurs. This system is found on somewhat excessively to excessively well-drained, deep sandy soils that are often associated with dune systems and ancient floodplains. In some areas, this system may actually occur as a result of overgrazing in Western Great Plains Tallgrass Prairie (CES303.673) or Western Great Plains Sand Prairie (CES303.670). Typically, this system is characterized by a sparse to moderately dense woody layer dominated by *Artemisia filifolia*, but other characteristic species may be present, including *Amorpha canescens*, *Prosopis glandulosa* (southern stands), *Prunus angustifolia*, *Prunus pumila* var. *besseyi* (northern stands), *Rhus trilobata*, and *Yucca glauca*. Associated herbaceous species can vary with geography, amount and season of precipitation, disturbance, and soil texture. The herbaceous layer typically has a moderate to dense canopy but may include stands with sparse understory. Several mid- to tallgrass species characteristic of sand substrates are usually present to dominant, such as *Andropogon hallii*, *Calamovilfa gigantea*, *Calamovilfa longifolia*, *Schizachyrium scoparium*, *Sporobolus cryptandrus*, *Sporobolus giganteus*, or *Hesperostipa comata*.

In the southern range of this system, *Quercus havardii* may also be present to dominant and represents one succession pathway that develops over time following a disturbance. *Quercus havardii* is able to resprout following a fire and thus may persist for long periods of time once established forming extensive clones. Edaphic and climatic factors are the most important dynamic processes for this type, with drought and extreme winds impacting this system significantly in some areas. Because *Quercus havardii* is able to resprout rapidly following fire, fire tends to cause structural changes in the vegetation, and compositional shifts are less significant in most cases. Overgrazing can lead to decreasing dominance of some of the grass species such as *Andropogon hallii*, *Calamovilfa gigantea*, and *Schizachyrium scoparium*. In the western extent of this system in the shortgrass prairie, more xeric mid- and shortgrass species such as *Hesperostipa comata*, *Sporobolus cryptandrus* and *Bouteloua gracilis* often dominate the herbaceous layer. This system is found primarily in semi-arid to arid areas of the Western Great Plains Division. It occurs on somewhat excessively to excessively well-drained and deep sandy soils. This system is often found associated with dune systems and/or ancient floodplains but may occur in soils derived from sandstone residuum.

HERBACEOUS

CES303.659 CENTRAL MIXEDGRASS PRAIRIE

This mixedgrass prairie system ranges from South Dakota into the Rolling Plains and the western Edwards Plateau of Texas. It is bordered by the shortgrass prairie on its western edge and the tallgrass prairie to the east. The loessal regions in west-central Kansas and central Nebraska, the Red Hills region of south-central Kansas and northern Oklahoma are all located within this system. Because of its proximity to other ecoregions, this system contains elements from both shortgrass and tallgrass prairies, which combine to form the mixedgrass prairie ecological system throughout its range. The distribution, species richness and productivity of plant species within the mixedgrass ecological system is controlled primarily by environmental conditions, in particular soil moisture and topography. Grazing and fire are important dynamic processes in this system. The relative dominance of the various grass and forb species within different associations in the system also can strongly depend on the degree of natural or human disturbance. This system can contain grass species such as *Bouteloua curtipendula*, *Schizachyrium scoparium*, *Andropogon gerardii*, *Hesperostipa comata*, *Sporobolus heterolepis*, and *Bouteloua gracilis*, although the majority of the associations within the region are dominated by *Pascopyrum smithii* or *Schizachyrium scoparium*. Numerous forb and sedge species (*Carex* spp.) can also occur within the mixedgrass system in the Western Great Plains. Although forbs do not always significantly contribute to the canopy, they can be very important. Some dominant forb species include *Ambrosia psilostachya*, *Echinacea angustifolia*, and *Lygodesmia juncea*. Oak species such as *Quercus macrocarpa* can occur also in areas protected from fire due to topographic position. This can cause an almost oak savanna situation in certain areas, although fire suppression may allow for a more closed canopy and expansion of bur oak beyond those sheltered areas. In those situations, further information will be needed to determine if those larger areas with a more closed canopy of bur oak should be considered part of Western Great Plains Dry Bur Oak Forest and Woodland (CES303.667). Likewise, within the mixedgrass system, small seeps may occur, especially during the wettest years. Although these are not considered a separate system, the suppression of fire within the region has enabled the invasion of both exotics and some shrub species such as *Juniperus virginiana* and also allowed for the establishment of *Pinus ponderosa* in some northern areas. This system is found primarily in the Central Mixed-grass Prairie (TNC Ecoregion 33); it becomes more restricted to mesic lowlands sites to the west and southwest in the shortgrass prairie region of Texas (S. Menard pers. comm. 2005). This is probably a reference to the Llano Estacado region rather than the Southern Shortgrass Prairie (TNC Ecoregion 28) (J. Teague pers. obs 2005). The Central Mixed-grass Prairie (TNC Ecoregion 33) should be extended south to include the Rolling Plains of Texas; being separated from the Southern Shortgrass Prairie (TNC Ecoregion 28) by the Caprock Escarpment (L. Elliott pers. comm. 2005).

Differences in topography and soil characteristics also occur across the range of this system. It is often characterized by rolling to extremely hilly landscapes with soils developed from loess, shale, limestone or sandstone parent material. Mollisol soils are most prevalent and range from silt loams and silty clay loams with sandy loams possible on the western edge of the range. The Red Hills region of Kansas and Oklahoma, which contains examples of this system, contains somewhat unique soil characteristics and has developed from a diversity of sources including red shale, red clay, sandy shale, siltstone, or sandstone. These soils have developed a characteristic reddish color from the primary material. These soils can consist of silt, loam, or clay and can have textures ranging from a fine sandy loam to a more clayey surface.

CES205.685 SOUTHEASTERN GREAT PLAINS TALLGRASS PRAIRIE

This system is found primarily within the Flint Hills and Osage Plains of Kansas and Oklahoma. Small patches can be found in the Ozarks of Missouri and the Arbuckle Mountains of Oklahoma. In southern Oklahoma and Texas, this is the primary natural system of the "Grand Prairie" or "Fort Worth Prairie," ranging south into the Lampasas Cutplain of Texas (EPA 29d and 29e, respectively). It is distinguished from Central Tallgrass Prairie (CES205.683) by having more species with southwestern geographic affinities and the presence of a thin soil layer over limestone beds ranging to more acidic substrates, although some areas of deeper soil are found within the region, especially on lower slopes, draws, and terraces. Because of the presence of the rocky substrate close to the surface and the rolling topography, this area is relatively unsuitable for agriculture. The Flint Hills contain one of the largest remaining, relatively intact pieces of tallgrass prairie. The vegetation in this system is typified by tallgrass species such as *Andropogon gerardii*, *Panicum virgatum*, *Schizachyrium scoparium*, and *Sorghastrum nutans* forming a dense cover. A moderate to high density of forb species also occurs. Species composition varies geographically, with *Oligoneuron rigidum* (= *Solidago rigida*), *Liatris punctata*, *Symphotrichum ericoides*, *Lespedeza capitata*, and *Viola pedatifida* occurring in some localities. Areas of deeper soil, especially lower slopes along draws, slopes and terraces, can include *Baptisia alba* var. *macrophylla*, *Liatris pycnostachya*, and *Vernonia missurica*. Shrub and tree species are relatively infrequent and, if present, constitute less than 10% cover in the area. Fire and grazing constitute the major dynamic processes for this region. Although many of the native common plant species still occur, grazing does impact this region. Poor grazing practices can lead to soil erosion and invasion by cool-season grasses such as *Bromus inermis* within its range.

This system is found primarily within the Flint Hills and Osage Plains of Kansas and Oklahoma. Small patches can be found in the Ozarks of Missouri and the Arbuckle Mountains of Oklahoma. In southern Oklahoma and Texas, this is the primary natural system of the "Grand Prairie" or "Fort Worth Prairie," ranging south into the Lampasas Cutplain of Texas (EPA 29d and 29e, respectively). In Missouri, it is attributed to EPA 40c, 40d, and possibly 39k.

This system is typified by the thin soil layer over limestone beds or acidic substrates such as chert or granite, although areas of deeper soils are possible along lower slopes, draws, and terraces. The topography is rolling and mostly unsuitable for agriculture.

CES205.684 TEXAS BLACKLAND TALLGRASS PRAIRIE

This system is found primarily in the Blackland Prairie region of Texas but can range into southern Oklahoma. It is typified by the presence of dark alkaline Vertisol soils over calcareous parent material interspersed with patches of acidic, sandy loam Alfisols and Mollisols. Microtopography such as gilgai and mima mounds can occur and are important microhabitats that lead to a high degree of plant diversity in this system. *Schizachyrium scoparium* and *Sorghastrum nutans* are the most frequent species with *Andropogon gerardii* as a possible associate, especially on the patches of Mollisol soils. *Tripsacum dactyloides* and *Panicum virgatum* are common associates on the Vertisol soils, especially on the gilgai microtopography. Fire and grazing constitute the major natural dynamics influencing this system. Infrequent, but intense, fires prevent woody species from establishing. Fire suppression and over grazing have allowed woody species to invade, and heavy grazing has allowed species such as *Buchloe dactyloides* and *Bouteloua rigidisetata* to invade.

This system is restricted to the Blackland Prairie region, part of the Crosstimbers and Southern Tallgrass Prairie Ecoregion, in Texas and possibly adjacent southern Oklahoma.

This system is typified by the presence of dark alkaline Vertisol soils over calcareous parent material interspersed with patches of acidic, sandy loam Alfisols and Mollisols. Microtopography such as gilgai and mima mounds can occur and are important microhabitats that lead to a high degree of plant diversity in this system.

The Main Belt of the Blackland Prairie is divided into Vertisol, Alfisol and Mollisol regions. The Vertisol region is characterized by the presence of dark clay alkaline soils over limestone marl parent material, while the Eastern Marginal prairies are characterized by variously textured Alfisols over sandstone parent material. Alkaline clay and clay loam Mollisols are found on the Austin Chalk formation on fragmented Cretaceous limestone. Two outlier prairies, the Fayette (EPA 32b) and San Antonio Prairies (EPA 33c), are underlain by both Vertisols and Alfisols. Each variation in soil texture and pH supports its characteristic community.

CES303.670 WESTERN GREAT PLAINS SAND PRAIRIE

The sand prairies constitute a very unique system within the western Great Plains. These sand prairies are often considered part of the tallgrass or mixedgrass regions in the western Great Plains but can contain elements from Western Great Plains Shortgrass Prairie (CES303.672), Central Mixedgrass Prairie (CES303.659), and Northwestern Great Plains Mixedgrass Prairie (CES303.674). The largest expanse of sand prairies (approximately 5 million ha) can be found in the Sandhills of north-central Nebraska and southwestern South Dakota. These areas are relatively intact. The primary use of this system has been grazing (not cultivation), and areas such as the Nebraska Sandhills can experience less degeneration than other prairie systems. Although greater than 90% of the Sandhills region is privately owned, the known fragility of the soils and the cautions used by ranchers to avoid poor grazing practices have allowed for fewer significant changes in the vegetation of the Sandhills compared to other grassland systems. The unifying and controlling feature for this system is that coarse-textured soils predominate and the dominant grasses are well-adapted to this condition. Soils in the sand prairies can be relatively undeveloped and are highly permeable. Soil texture and drainage along with a species' rooting morphology, photosynthetic physiology, and mechanisms to avoid transpiration loss are highly important in determining the composition of the sand prairies. In the northwestern portion of its range, stand size corresponds to the area of exposed caprock sandstone, and small patches predominate, but large patches are also found embedded in the encompassing Northwestern Great Plains Mixedgrass Prairie (CES303.674). Another important feature is their susceptibility to wind erosion. Blowouts and sand draws are some of the unique wind-driven disturbances in the sand prairies, particularly the Nebraska Sandhills. In most of eastern Montana, substrates supporting this system have weathered in place from sandstone caprock; thus the solum is relatively thin, and the wind-sculpted features present further east, particularly in Nebraska, do not develop. Graminoid species dominate the sand prairies, although relative dominance can change due to impacts of wind disturbance. *Andropogon hallii* and *Calamovilfa longifolia* are the most common species, but other grass and forb species such as *Hesperostipa comata*, *Carex inops ssp. heliophila*, and *Panicum virgatum* may be present. Apparently only *Calamovilfa longifolia* functions as a dominant throughout the range of the system. In the western extent, *Hesperostipa comata* becomes more dominant, and *Andropogon hallii* is less abundant but still present. Communities of *Artemisia cana ssp. cana* are included here in central and eastern Montana. Patches of *Quercus havardii* can also occur within this system in the southern Great Plains. Fire and grazing constitute the other major dynamic processes that can influence this system. This system was edited to expand the concept to include sandy portions of the mixedgrass prairie of the Montana plains. Although in terms of potentially dominant

graminoids there is virtually a complete overlap between the eastern and western extremities of the system, there is a distinct shift from west to east from midgrass species dominance, most notably *Hesperostipa comata*, to tallgrass species dominance, including prominently *Andropogon gerardii* and *Andropogon hallii*. Prevailing patch size also shifts from smaller to larger moving west to east. Current thinking is to include this variation within this system, but with more information and input from other Great Plains ecologists in the U.S. and Canada, this concept is subject to change, including the possibility of creating a new system.

This system is found throughout the Western Great Plains Division. The largest and most intact example of this system is found within the Sandhills region of Nebraska and South Dakota. However, it is also common (though occurring in predominantly small patches) farther west into central and eastern Montana. Its western extent in Wyoming is still to be determined, but it does occur in mapzone 29 on weathered-in-place sandy soils, where *Calamovilfa longifolia* is found, along with *Artemisia cana*.

The distribution, species richness and productivity of plant species within the sand prairie ecological system are controlled primarily by environmental conditions, in particular the temporal and spatial distribution of soil moisture and topography. Soils in the sand prairies can be relatively undeveloped and are highly permeable. Soil texture and drainage along with a species' rooting morphology, photosynthetic physiology, and mechanisms to avoid transpiration loss are highly important in determining the composition and distribution of communities/associations within the sand prairies. Another important aspect of soils in the sand prairies is their susceptibility to wind erosion. Blowouts and sand draws are some of the unique wind-driven disturbances in the sand prairies, particularly the Nebraska Sandhills, which can profoundly impact vegetation composition and succession within this system. This tallgrass system is found primarily on sandy and sandy loam soils that can be relatively undeveloped and highly permeable as compared to Western Great Plains Tallgrass Prairie (CES303.673), which occurs on deeper loams. This system is usually found in areas with a rolling topography and can occur on ridges, midslopes and/or lowland areas within a region. It often occurs on moving sand dunes, especially within the Sandhills region of Nebraska and South Dakota. In Montana, occurrences are intimately associated with Northwestern Great Plains Mixedgrass Prairie (CES303.674), usually occupying higher positions in local landscapes due to the fact that sandy members of some formations (that are predominantly marine shales) constitute the highest (and most weathering-resistant) points in the landscape.

WOODY WETLAND

CES303.651 EDWARDS PLATEAU FLOODPLAIN

This system occurs on floodplain terraces along perennial rivers and streams in central Texas. Canopy dominants may include *Ulmus crassifolia*, *Juniperus ashei*, *Celtis laevigata*, *Quercus fusiformis*, *Fraxinus texensis*, *Platanus occidentalis*, *Acer negundo*, *Juglans major*, *Quercus macrocarpa*, or *Carya illinoensis*. *Carya illinoensis* may be more likely to occur in deeper and better-developed alluvial soils. Apparent dominance of *Carya illinoensis* may also be an artifact of preferential harvesting of other species, leaving this species in greater abundance. Alluvial sedimentation processes dominate the formation and maintenance of this system. However, overgrazing and/or overbrowsing may influence recruitment of overstory species and composition of the understory and herbaceous layers. This system occurs along larger permanent rivers and streams throughout the Edwards Plateau of Texas and possibly adjacent ecoregions. It occurs from the Leon watershed in the Limestone Cutplain (EPA 29e) south to the edge of the Bacones Canyonlands (EPA 30c), west through the Edwards Plateau and north to the Pecan Bayou and Concho River watersheds in the lower Limestone Plains (EPA 27j) and lower Crosstimbers (EPA 29c) (EPA 2001). This system occurs on alluvial terraces along permanent rivers and streams in central Texas.

CES303.652 EDWARDS PLATEAU RIPARIAN

This system occurs in various situations along small and intermittent streams of the Edwards Plateau, with drier representatives occurring in the western plateau and the Stockton Plateau, and moister representatives (such as communities dominated by *Juglans microcarpa* and *Brickellia laciniata*) in the eastern plateau. Representatives of this system typically occur in stream-scoured situations and vary in the openness of the habitat and physiognomy.

CES205.710 SOUTHEASTERN GREAT PLAINS FLOODPLAIN FOREST

This ecological system is found in the floodplains of medium and larger rivers of the East Central Texas Plains, Texas Blackland Prairie Regions, Crosstimbers, and the southeastern edge of the Central Great Plains (Level 3 Ecoregions 33, 32, 29 and 27 respectively, *sensu* Griffith et al. (2004)). Alluvial soils and sedimentation processes typify this system. Periodic, intermediate flooding and deposition (every 5-25 years) dominates the formation and maintenance of this system. Dominant communities within this system range from floodplain forests to wet meadows to gravel/sand flats; however, they are linked by underlying soils and the flooding regime. Canopy dominants may include *Carya illinoensis*, *Ulmus crassifolia*, *Ulmus americana*, *Celtis laevigata*, *Quercus nigra*, *Platanus occidentalis*, *Acer negundo*, *Quercus macrocarpa*, *Morus rubra*, *Fraxinus pennsylvanica*, *Salix nigra*, and *Sapindus saponaria* var. *drummondii* (= *Sapindus drummondii*). Overgrazing and/or overbrowsing may influence recruitment of overstory species and composition of the understory and herbaceous layers. Shrub species may include *Callicarpa americana*, *Ilex decidua*, *Ilex americana*, *Sideroxylon lanuginosum*, *Diospyros virginiana*, *Juniperus virginiana*, *Cornus drummondii*, and *Viburnum rufidulum*, which may occur as dense patches following disturbance, but are otherwise generally fairly sparse. Vines such as *Berchemia scandens*, *Campsis radicans*, *Vitis* spp., *Parthenocissus quinquefolia*, and *Ampelopsis arborea* may be conspicuous. Herbaceous cover includes *Elymus virginicus*, *Verbesina virginica*, *Chasmanthium latifolium*, *Chasmanthium sessiliflorum*, *Tripsacum dactyloides*, *Symphotrichum drummondii* var. *texanum*, *Geum canadense*, *Sanicula canadensis*, *Panicum virgatum*, *Galium* spp., and *Carex* sp. Herbaceous cover may be quite high, especially in situations where shrub cover is low. The environment and vegetation of this system become generally and correspondingly drier from east to west with moister representatives (such as communities containing *Quercus phellos*, *Quercus pagoda*, *Quercus alba*, and *Quercus lyrata*) occurring along the eastern and northeastern margins of the range. Representatives of this system may vary in the openness of the habitat and physiognomy.

This system occupies relatively broad flats at low topographic positions, along large streams where alluvial deposition dominates. It is found in the floodplains of medium and larger rivers of the East Central Texas Plains, Texas Blackland Prairie Regions, Crosstimbers, and the southeastern edge of the Central Great Plains (Level 3 Ecoregions 33, 32, 29 and 27 respectively, *sensu* Griffith et al. (2004)). Soils are primarily alluvial and range from sandy to dense clays.

CES205.709 SOUTHEASTERN GREAT PLAINS RIPARIAN FOREST

This ecological system occurs in various situations along small and intermittent streams in the East Central Texas Plains, Texas Blackland Prairie Regions, Crosstimbers, and the southeastern edge of the Central Great Plains (Level 3 Ecoregions 33, 32, 29 and 27, respectively, *sensu* Griffith et al. (2004)). Some trees that may be present in stands of this system include *Celtis laevigata* var. *laevigata*, *Celtis laevigata* var. *reticulata*, *Platanus occidentalis*, *Quercus nigra*, *Quercus phellos*, *Amorpha fruticosa*, *Forestiera*

acuminata, *Acer saccharinum*, *Sapindus saponaria*, *Salix nigra*, *Fraxinus pennsylvanica*, *Gleditsia triacanthos*, *Carya illinoensis*, and *Ulmus crassifolia*. The environment and vegetation of this system become generally and correspondingly drier from east to west with moister representatives (such as communities containing *Quercus nigra*) occurring in the eastern parts of the range. Representatives of this system typically occur in stream-scoured situations and vary in the openness of the habitat and physiognomy.

This system occurs on minor intermittent streams and tributaries throughout the East Central Texas Plains, Texas Blackland Prairie Regions, Crosstimbers, and the southeastern edge of the Central Great Plains (Level 3 Ecoregions 33, 32, 29 and 27 respectively, sensu Griffith et al. (2004)). It is found along medium to very small, intermittent to ephemeral drainages. This type is ubiquitous throughout, but species composition and flood regimes are variable and are thought to be dependent on soil and geologic substrates. Generally, these are less thick alluvium than in floodplain terraces. These are flashy streams, and flooding rather than fire will be the dominant process in this system. Fuels in this system are variable, and fire-return interval is partially determined by that of the adjacent and surrounding matrix upland system, where fuels are present.

BARREN

CES203.398 SOUTHEASTERN COASTAL PLAIN CLIFF

This ecological system consists of steep to vertical or overhanging outcrops of unconsolidated sediment or rock in the Gulf and Atlantic Coastal Plain. They occur on lower bluffs adjacent to rivers or streams. The vegetation is generally sparse, limited to plants growing on bare substrate, small ledges, and other favorable microsites. The flora is a mix of herbs, shrubs, vines, tree seedlings, and some larger trees, with bryophytes potentially dominant in some examples. It typically includes opportunistic species of open and disturbed areas, along with species from adjacent forest communities and other species of wet and dry open areas. Dense shrubby or woodland vegetation may occur on the edges. Occasional examples may have denser bryophyte vegetation. More information is needed on the associations that belong to this system. Compared to cliffs of other regions, the soft Coastal Plain cliffs tend to be more frequently disturbed and more dynamic. Soil development and primary succession would quickly replace any examples that are not periodically disturbed and renewed by slumping, generally caused by undercutting by streams. Smaller slumps and active erosion by upland runoff may also disturb the communities. This is the primary rock outcrop system of the Coastal Plain, but small shaded outcrops of limestone or sandstone may be included in other systems.

These cliffs occur on steep to vertical or overhanging outcrops of Coastal Plain substrates, usually sand, clay, or sandstone. Outcrops typically occur along rivers or streams, where undercutting causes slumps that periodically renew the bare substrate. Most of these cliffs are dry, but small zones of seepage are often present, especially at the top. One association represents small clay cliffs that are kept wet by seepage. Lower portions of cliffs may periodically flood.